

BIOMIN[®] BOOSTER V



PRODUCT INFORMATION

BIOMIN BOOSTER V is a chelated multi-mineral which is chelated mainly by Glycine and other amino acids.

BIOMIN BOOSTER V was designed for soil and/or foliar applications. Completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. Designed to prevent and correct trace mineral deficiencies and to boost crops during critical or fast growing periods.

BIOMIN BOOSTER V can be applied to ALL crops including all vegetables, potatoes, carrots, onions, all berries and strawberries, all brassicas, all fruit trees, avocados, persimmons, citrus, all nut trees, vineyards, kiwifruit, tropical fruit, lettuce, asparagus, all melons, beans, ginger, ornamentals, cut flowers and pastures.

GUARANTEED ANALYSIS

As true amino acid Chelates:	Dry Weight Basis %w/w
Magnesium (Mg)	4.0%
Zinc (Zn)	4.0%
Iron (Fe)	2.6%
Manganese (Mn)	2.6%
Copper (Cu)	1.5%
Boron (B)	1.5%
Molybdenum (Mo)	0.5%
As amino acid:	
Nitrogen (N)	4%

Magnesium, Zinc, Iron & Manganese present as Glycine chelates. Molybdenum & Boron present as Amino Acid complexes. Nitrogen present as Amino Acid.

COMPATIBILITY

Always perform a compatibility test before mixing any product with **BIOMIN BOOSTER V**. Not compatible with Copper or Copper containing products and care should be taken with crops that have had Copper applied.

PREVENTS AND CORRECTS MINERAL DEFICIENCIES.

In conditions where the water and/or soil pH is alkaline, it would be advantageous to apply **MOBILIZER** to the tank water before adding **BIOMIN BOOSTER V**. To further increase availability, add **BIOMIN BOOSTER V** to the tank when the pH of the water is 6.5 - 7.5.

WARNING

Harmful if swallowed.
Causes eye irritation.
Causes skin irritation.
Suspected of damaging fertility or the unborn child.
May cause damage to organs through prolonged or repeated exposure.
Harmful to aquatic life. May cause long lasting harmful effects to aquatic life.



PRECAUTIONS

Keep out of reach of children. Read label before use. Do not handle until all safety precautions have been read and understood. Avoid breathing dust. Wash hands thoroughly after handling. Do not eat, drink, or smoke when using this product. Wear protective gloves/clothing. Avoid release to the environment.

Wash contaminated clothing before reuse. If exposed or concerned: Get medical advice if you feel unwell.

IF SWALLOWED: Rinse mouth. Call a POISON CENTRE or doctor if you feel unwell.

IF ON SKIN: Wash with plenty of water. If skin irritation occurs: Get medical advice.

IF IN EYES: Rinse cautiously with water for several minutes. If irritation persists: Get medical advice.

Store locked up. Dispose of contents/container in accordance with local regulations.

24-hour emergency number: 0800 764 766 (NZ Poisons Centre)

Approved Pursuant to the HSNO Act 1996, under the Fertilisers – Subsidiary Hazard group standard HSR002571. See www.epa.govt.nz for approval conditions.



**Net Contents: 10kg
Lot#**

Best Before Date:

Imported and distributed in New Zealand exclusively by: Roots, Shoots & Fruits Ltd, C/- 27 Hastie Avenue, Auckland, New Zealand. Ph: +64 (0) 9372 9155 Email: rsf@rd2.co.nz
Website: www.rd2.co.nz Manufactured by: JH BIOTECH, INC. PO Box 3538, Ventura, CA 93006, USA. www.jhbiotech.com This product embodies proprietary or patented information and may be used only for agricultural purposes. Any duplication, reproduction, breeding, transformation or reverse engineering is prohibited.

SOIL HEALTH · PLANT HEALTH · YOUR HEALTH

BIOMIN[®] BOOSTER V



Directions and General Recommendations

SOIL APPLICATION

BIOMIN BOOSTER V may be applied before or at planting through irrigation water. The application rate is 1.5 - 3 kilograms per hectare depending on crop, soil conditions and application methods.

FOLIAR APPLICATION

ALL FIELD CROPS: (grain crops, flowers, etc...): Apply 1 - 1.5 kilograms of **BIOMIN BOOSTER V** per hectare when crops are 15 - 30cms high or when deficiency symptoms appear. Repeat application 2 - 4 weeks later or as needed.

ALL VEGETABLE CROPS: Apply 1 - 1.5 kilograms of **BIOMIN BOOSTER V** per hectare at about 2 - 4 weeks after emergence or if deficiency symptoms appear. Repeat the application every 2 - 3 weeks or as needed.

GREENHOUSE CROPS: Apply 100 - 150 grams of **BIOMIN BOOSTER V** per 100 litres of water 2-3 weeks after emergence or transplanting, or when trace element deficiencies first appear. Repeat the application every month.

APPLES: Apply 1 - 2 kilograms of **BIOMIN BOOSTER V** per hectare as a post harvest application only at least 6 weeks before dormancy.

FRUIT TREES: Apply 1 - 2 kilograms of **BIOMIN BOOSTER V** per hectare as a post harvest application only at least 6 weeks before dormancy.

KIWIFRUIT: Apply 1 - 1.5 kilograms of **BIOMIN BOOSTER V** per hectare in early Spring when new leaves emerge or 2 weeks before flowering. Repeat the application postharvest at 6 weeks before dormancy at a rate of 1 - 2 kilograms per hectare.

GRAPES & CITRUS: Apply 1 - 1.5 kilograms of **BIOMIN BOOSTER V** per hectare in early Spring when new leaves emerge or 2 weeks before flowering or at any stage in the season when deficiency symptoms appear. Repeat the application postharvest at 6 weeks before dormancy at a rate of 1 - 2 kilograms per hectare.

BIOMIN BOOSTER V is best applied when plants are in an active growing stage, after irrigation or rainfall. For foliar application, it should be diluted with at least 100 times of water.

BIOMIN BOOSTER V may be used at any time throughout the season. For best results and more specific recommendations, contact Roots, Shoots & Fruits.

LIMITED WARRANTY

Manufacturer or reseller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor reseller shall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label.

Imported and distributed in New Zealand exclusively by: Roots, Shoots & Fruits Ltd, C/- 27 Hastie Avenue, Auckland, New Zealand. Ph: +64 (0) 9372 9155 Email: rsf@rd2.co.nz
Website: www.rd2.co.nz Manufactured by: JH BIOTECH, INC. PO Box 3538, Ventura, CA 93006, USA. www.jhbiotech.com This product embodies proprietary or patented information and may be used only for agricultural purposes. Any duplication, reproduction, breeding, transformation or reverse engineering is prohibited.

SOIL HEALTH · PLANT HEALTH · YOUR HEALTH